



**Fermi National Accelerator Laboratory**

**FERMILAB-Pub-96/393-E**

**E704**

**Spin Transfer in Inclusive  $\Lambda^0$  Production by Transversely Polarized  
Protons at 200 GeV/c**

A. Bravar et al.

The E704 Collaboration

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

September 1997

Submitted to *Physical Review Letters*

## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

## **Distribution**

*Approved for public release; further dissemination unlimited.*

# Spin Transfer in Inclusive $\Lambda^0$ Production by Transversely Polarized Protons at 200 GeV/c

A. Bravar,<sup>17,\*</sup> D.L. Adams,<sup>16</sup> N. Akchurin,<sup>6</sup> N.I. Belikov,<sup>5</sup> B.E. Bonner,<sup>16</sup> J. Bystricky,<sup>2</sup>  
M.D. Corcoran,<sup>16</sup> J.D. Cossairt,<sup>3</sup> J. Cranshaw,<sup>16,†</sup> A.A. Derevschikov,<sup>5</sup> H. En'yo,<sup>8</sup>  
H. Funahashi,<sup>8</sup> Y. Goto,<sup>8,‡</sup> O.A. Grachov,<sup>5</sup> D.P. Grosnick,<sup>1</sup> D.A. Hill,<sup>1</sup> T. Iijima,<sup>8</sup>  
K. Imai,<sup>8</sup> Y. Itow,<sup>8,§</sup> K. Iwatani,<sup>4</sup> K. Kuroda,<sup>11</sup> M. Laghai,<sup>1</sup> F. Lehar,<sup>2</sup> A. de Lesquen,<sup>2</sup>  
D. Lopiano,<sup>1</sup> F.C. Luehring,<sup>13,\*\*</sup> T. Maki,<sup>7</sup> S. Makino,<sup>8,††</sup> A. Masaike,<sup>8</sup> Yu.A. Matulenko,<sup>5</sup>  
A.P. Meschanin,<sup>5</sup> A. Michalowicz,<sup>11</sup> D.H. Miller,<sup>13</sup> K. Miyake,<sup>8</sup> T. Nagamine,<sup>8</sup>  
F. Nessi-Tedaldi,<sup>16,‡‡</sup> M. Nessi,<sup>16,‡‡</sup> C. Nguyen,<sup>16</sup> S.B. Nurushev,<sup>5</sup> Y. Ohashi,<sup>1</sup> Y. Onel,<sup>6</sup>  
D.I. Patalakha,<sup>5</sup> G. Pauletta,<sup>18</sup> A. Penzo,<sup>17</sup> G.F. Rappazzo,<sup>12</sup> A.L. Read,<sup>3</sup> J.B. Roberts,<sup>16</sup>  
L. van Rossum,<sup>2</sup> V.L. Rykov,<sup>5</sup> N. Saito,<sup>8,‡</sup> G. Salvato,<sup>12</sup> P. Schiavon,<sup>17</sup> J. Skeens,<sup>16</sup>  
V.L. Solovyanov,<sup>5</sup> H. Spinka,<sup>1</sup> R.W. Stanek,<sup>1</sup> R. Takashima,<sup>9</sup> F. Takeutchi,<sup>10</sup> N. Tamura,<sup>14</sup>  
D.G. Underwood,<sup>1</sup> A.N. Vasiliev,<sup>5</sup> J.L. White,<sup>16,§§</sup> S. Yamashita,<sup>8</sup> A. Yokosawa,<sup>1</sup>  
T. Yoshida,<sup>15</sup> and A.M. Zanetti<sup>17</sup>

(Fermilab E704 Collaboration)

<sup>1</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*

<sup>2</sup>*CEA-DAPNIA/SPP, CE-Saclay, F-91191 Gif-sur Yvette, France*

<sup>3</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

<sup>4</sup>*Hiroshima University, Higashi-Hiroshima 724, Japan*

<sup>5</sup>*Institute of High Energy Physics, 142284 Protvino, Russia*

<sup>6</sup>*Department of Physics and Astronomy, University of Iowa, Iowa City, Iowa 52242, USA*

<sup>7</sup>*University of Occupational and Environmental Health, Kita-Kyushu 807, Japan*

<sup>8</sup>*Department of Physics, Kyoto University, Kyoto 606-01, Japan*

<sup>9</sup>*Kyoto University of Education, Kyoto 612, Japan*

<sup>10</sup>*Kyoto-Sangyo University, Kyoto 612, Japan*

<sup>11</sup>*Laboratoire de Physique des Particules, BP 909, 74017 Annecy-le-Vieux, France*

<sup>12</sup>*Dipartimento di Fisica, Università di Messina and INFN Messina, I-98100 Messina, Italy*

<sup>13</sup>*Physics Department, Northwestern University, Evanston, Illinois 60201, USA*

<sup>14</sup>*Department of Physics, Okayama University, Okayama 800, Japan*

<sup>15</sup>*Osaka City University, Osaka 558, Japan*

<sup>16</sup>*T.W. Bonner Nuclear Laboratory, Rice University, Houston, Texas 77251, USA*

<sup>17</sup>*Dipartimento di Fisica, Università di Trieste and INFN Trieste, I-34100 Trieste, Italy*

<sup>18</sup>*Università di Udine and INFN Udine, I-33100 Udine, Italy*

---

\*Universität Mainz, D-55099 Mainz, Germany

†INFN Trieste, I-34100 Trieste, Italy

‡RIKEN, Saitama 351-01, Japan

§Inst. for Cosmic Ray Research, University of Tokyo, Gifu 506-12, Japan

\*\*Indiana University, Bloomington, IN 47405, USA

††Wakayama Medical College, Wakayama 649-63, Japan

‡‡CERN, CH-1211 Geneva 23, Switzerland

§§SLAC, Stanford, CA 94309, USA

# Abstract

Surprisingly large polarizations in hyperon production have been known for a long time. The spin dynamics of the production process can be further investigated with polarized beams. Recently, a negative asymmetry  $A_N$  was found in inclusive  $\Lambda^0$  production with a 200 GeV/ $c$  transversely polarized proton beam. The depolarization  $D_{NN}$  in  $p \uparrow + p \rightarrow \Lambda^0 + X$  has been measured with the same beam over a wide  $x_F$  range and at moderate  $p_T$ .  $D_{NN}$  reaches positive values of about 30 % at high  $x_F$  and  $p_T \sim 1.0$  GeV/ $c$ . This result shows a sizeable spin transfer from the incident polarized proton to the outgoing  $\Lambda^0$ .

PACS numbers: 13.88.+e, 13.85.Ni, 14.20.Jn

The observation 20 years ago of a large negative polarization in inclusive  $\Lambda^0$  production by an unpolarized proton beam at 300 GeV/c [1] renewed interest in spin as an important factor in high-energy hadron interactions. Afterwards several experiments measured large polarizations for various hyperons over a wide kinematical range [2]. Previous expectations, based on Regge theory and QCD predictions, were that spin effects would vanish at high energies, since the smallness of spin-flip amplitudes and the contribution of several production channels to an inclusive process with large multiplicity of final states make it unlikely to have the coherent interference between spin non-flip and spin flip amplitudes that leads to sizeable polarization effects. Recently, a significant negative analyzing power  $A_N$  has been found at 200 GeV/c in inclusive  $\Lambda^0$  production by a transversely polarized proton beam at high  $x_F$  and moderate  $p_T$  ( $p_T \sim 1$  GeV/c) [3]. Large  $A_N$  values have been also found in inclusive pion production with the same proton and antiproton polarized beam [4,5].

Different quark-parton models using static SU(6) wave functions were proposed to interpret these polarization effects by introducing a spin dependence into the partonic fragmentation and recombination processes [6–8]. The  $\Lambda^0$  polarization is attributed to some mechanism, based on semiclassical arguments [6,7] or inspired to QCD [8], by which *strange* quarks produced in the fragmentation process acquire a large negative polarization. The features of the pion  $A_N$  data [4,5] are compatible with these models, provided that this effect occurs also for *up* and *down* quarks. The spin dynamics of these processes can be further investigated using polarized proton beams. In the previous models no correlation with the incident proton polarization is expected in inclusive  $\Lambda^0$  production, since the  $\Lambda^0$  spin is carried entirely by its constituent *strange* quark and the *ud* di-quark (which is in a spin and isospin singlet state) propagates unperturbed as a spectator in the interaction. Therefore spin asymmetries related to the beam polarization are expected to vanish. The negative asymmetry  $A_N$  observed in  $\Lambda^0$  production [3] is difficult to integrate in this picture unless the spectator *ud* di-quarks play a more significant role in the recombination process than generally expected [9]. Studies of other spin asymmetries in high energy hyperon production add further input to understand these phenomena.

In this Letter we report on the measurement of the depolarization parameter  $D_{NN}$  in inclusive  $\Lambda^0$  production with the 200 GeV/c Fermilab polarized proton beam [10] and a 1.0 meter long liquid hydrogen target in the kinematical range  $0.2 \leq x_F \leq 1.0$  and  $0.1 \leq p_T \leq 1.5$  GeV/c. The double-spin parameter

$$D_{NN} = \frac{E \frac{d^3\sigma}{dp^3}^{\uparrow\uparrow} - E \frac{d^3\sigma}{dp^3}^{\uparrow\downarrow}}{E \frac{d^3\sigma}{dp^3}^{\uparrow\uparrow} + E \frac{d^3\sigma}{dp^3}^{\uparrow\downarrow}}$$

measures the fraction of the incident proton polarization transferred to the inclusively produced  $\Lambda^0$ .  $E \frac{d^3\sigma}{dp^3}^{\uparrow\uparrow(\uparrow\downarrow)}$  is the spin-dependent differential cross-section for the process  $p \uparrow + p \rightarrow \Lambda^0 \uparrow + X$  with parallel (anti-parallel) spin configurations for the incident proton and the outgoing  $\Lambda^0$ , both polarizations being orthogonal to the production plane.

The transversely polarized proton beam contained simultaneously protons of opposite *tagged* polarizations. This considerably suppressed systematic effects. The average beam polarization was  $0.46 \pm 0.03$  for both signs. Typical beam intensities at the experimental target were of the order of  $2 \times 10^7$  polarized protons per 20 second spill.

$\Lambda^0$  hyperons produced at the experimental target were identified by reconstructing their decay  $\Lambda^0 \rightarrow p\pi^-$ . Charged particles were measured in a forward spectrometer, described in Refs. [3,4], equipped with 42 multi-wire proportional planes and a 3 T-m  $\int Bdl$  dipole analyzing magnet. A threshold Cherenkov counter, C1, downstream of the magnet, was used for proton identification.

Secondary  $\Lambda^0$  decay vertices ( $V^0$ 's) were searched by combining proton tracks identified by C1 with negatively charged tracks assuming that they were  $\pi^-$ 's [3]. It was required that: (1) the closest distance in space of the two tracks was  $< 2$  mm and that the  $V^0$  decay vertex was located between 20 cm and 540 cm downstream of the target end; (2) the  $V^0$  came from the target and matched there the beam impact point within 2 mm in the transverse plane; (3) the  $V^0$ 's populated the decay phase space region corresponding to  $\Lambda^0$  decays, bounded by  $0.45 < (p_L^+ - p_L^-)/(p_L^+ + p_L^-) < 0.95$  and  $q_T = q_T^+ = q_T^- < 0.12$  GeV/c, where  $p_L^+$  ( $p_L^-$ ) is the longitudinal component of the positive (negative) track momentum and  $q_T$  is the transverse component of these momenta with respect to the  $V^0$  line of flight.

These selection cuts led to a clean  $\Lambda^0$  peak in the  $p\pi^-$  invariant mass spectrum, centered at  $1.116 \text{ GeV}/c^2$  with a width  $\sigma = 1.7 \text{ MeV}/c^2$  and an estimated 2 % uniform background below the  $\Lambda^0$  peak, as most sources of background were suppressed by the vertex fiducial volume cut, the selection of the  $\Lambda^0$  decay phase space region, and  $K_S^0 \rightarrow \pi^+\pi^-$  decays were rejected by C1. We selected  $\Lambda^0$ 's in a mass window of  $\pm 5.1 \text{ MeV}/c^2$  about the peak. For the  $D_{NN}$  analysis we required additionally that the  $\Lambda^0$ 's were produced to the beam right in the azimuthal angular interval of  $\pm 60^\circ$  from the horizontal plane around the beam axis. A sample of about 40,000  $\Lambda^0$ 's was thus selected.

For each  $x_F$  and/or  $p_T$  bin, the double-spin parameter  $D_{NN}$  was extracted from the  $\Lambda^0$  decay proton angular distribution in the  $\Lambda^0$  rest frame by defining 4 sets of events, integrating the decay proton angular distribution above and below the  $\Lambda^0$  production plane for the two opposite beam polarizations separately.  $D_{NN}$  is then obtained from the asymmetry:

$$D_{NN} = \frac{1}{P_B \langle \cos \phi_V \rangle} \frac{2}{\alpha_\Lambda} \frac{(N_{up}^+ + N_{down}^-) - (N_{up}^- + N_{down}^+)}{(N_{up}^+ + N_{down}^-) + (N_{up}^- + N_{down}^+)}$$

where, for instance,  $N_{up}^+$  is the number of  $\Lambda^0$ 's produced by beam protons polarized upward and emitting decay protons in the positive hemisphere with respect to the normal to the production plane.  $P_B$  is the proton beam polarization,  $\phi_V$  is the angle between the beam polarization axis directed upward and the normal to the production plane ( $\langle \cos \phi_V \rangle \approx -0.85$  in the selected azimuthal range), and  $\alpha_\Lambda = 0.642$  is the  $\Lambda^0$  decay asymmetry.

The  $D_{NN}$  results thus obtained, are independent, to a good accuracy, of apparatus and reconstruction biases, since  $\Lambda^0$ 's were measured with the same apparatus and opposite beam polarizations simultaneously. For a check of systematic biases we evaluated  $D_{NN}$  for non- $\Lambda^0$  events ( $p\pi^-$  combinations outside of the  $\Lambda^0$  mass window and  $K_S^0$ ), and we found that  $D_{NN} = 0.007 \pm 0.039$  for these background events. We also evaluated the  $\Lambda^0$  polarization by averaging over opposite beam polarizations and found a good agreement with existing polarization results.

The depolarization  $D_{NN}$  is given in Table 1 and shown in Figure 1 as a function of  $p_T$  averaged over the  $x_F$  interval of 0.2–1.0.  $D_{NN}$  increases with  $p_T$  to significantly large



positive values with an indication of flattening in  $p_T$  above 1.0 GeV/ $c$ , while at low  $p_T$  values it is compatible with zero. Figure 2 and Table 2 show the double-spin parameter  $D_{NN}$  as a function of  $x_F$  averaged over the  $p_T$  interval of 0.1–1.5 GeV/ $c$ . At large  $x_F$  values  $D_{NN}$  reaches positive values as large as 30 % at  $p_T \sim 1$  GeV/ $c$  (see also Table 1 rows 6–11, where the  $p_T$  dependence is shown for two separate  $x_F$  intervals), while almost no dependence in  $x_F$  is observed for  $x_F < 0.6$ , where  $D_{NN}$  is compatible with zero or slightly positive. In Figure 3 the  $D_{NN}$  data are split into two  $p_T$  intervals and plotted as a function of  $x_F$ . At low  $p_T$  they appear to be essentially zero, while in the high  $p_T$  bin they show large positive values. A few  $D_{NN}$  measurements in inclusive  $\Lambda^0$  production were previously performed with polarized proton beams at much lower energies of 6 GeV/ $c$  [11], 13.3 GeV/ $c$  and 18.5 GeV/ $c$  [12]. Figure 2 shows also data obtained at 18.5 GeV/ $c$  for a bin about  $\langle p_T \rangle \sim 1.0$  GeV/ $c$  [12]. These data appear to be compatible with zero over the measured range, which however doesn't extend above  $x_F \sim 0.5$ . More recently, a sizeable spin transfer has been inferred in  $\Omega^-$  production by high energy neutral beams containing transversely polarized  $\Lambda^0$ 's and  $\Xi^0$ 's [13].

The kinematical dependence of present  $D_{NN}$  results shows, in magnitude, a behavior similar to the hyperon polarization [2]. The observed  $D_{NN}$  results, as in the case of the  $A_N$  data in inclusive  $\Lambda^0$  production [3], cannot however be directly obtained from a mechanism such as proposed to explain the  $\Lambda^0$  polarization [6–8], where a highly polarized *strange* quark produced in the fragmentation process recombines with an unpolarized *ud* spectator di-quark from the incident proton independently of its polarization. Our results indicate a substantial spin transfer from the incident polarized proton to the inclusively produced  $\Lambda^0$  as large as 30 % at high  $x_F$  ( $x_F > 0.6$ ) and  $p_T \sim 1$  GeV/ $c$ .

More recent models based on non-perturbative approaches and peripheral mechanisms with an underlying quasi-binary subprocess, such as a  $\pi$  exchange mechanism [14], or resonance-decay interference between real and virtual channels [15], were proposed to explain the  $\Lambda^0$  polarization. These models might also accomodate a more substantial spin dependence in the  $\Lambda^0$  production process, such as the one shown by the present data. A

model, based on the idea of rotating constituents in polarized protons [16], is fairly successful in accounting for the observed  $A_N$  behavior in pion production. This model appears to reproduce qualitatively also the  $\Lambda^0$  analyzing power and the  $D_{NN}$  data presented in this Letter.

We are grateful for the assistance of the staff of Fermilab and of all the participating institutions. This work would not have been possible without the ideas and efforts of our colleagues M.M. Gazzaly, R. Rzaev, N. Tanaka, and A. Villari; we regret their passing away and we will miss them all greatly. This experiment was supported in part by the U.S. Department of Energy, the Istituto di Fisica Nucleare in Italy, the Commissariat à l'Energie Atomique and the Institut National de Physique Nucléaire et de Physique des Particules in France, the former U.S.S.R. Ministry of Atomic Power and Industry, and the Ministry of Education Science and Culture in Japan.

## REFERENCES

- [1] G. Bunce *et al.*, Phys. Rev. Lett. **36**, 1113 (1976).
- [2] for a review see L.G. Pondrom, Phys. Rep. **122C**, 57 (1985); K. Heller, in *Proceedings of the 9th Symposium on High Energy Spin Physics, Bonn 1990*, edited by K.H. Althoff and W. Meyer (Springer-Verlag, Berlin 1991); and references therein.
- [3] A. Bravar *et al.*, Phys. Rev. Lett. **75**, 3073 (1995).
- [4] A. Bravar *et al.*, Phys. Rev. Lett. **77**, 2028 (1996); D.L. Adams *et al.*, Phys. Lett. **264B**, 462 (1991); D.L. Adams *et al.*, Phys. Lett. **261B**, 201 (1991).
- [5] for a review see A. Bravar, in *Proceedings of the Adriatico Research Conference on Trends in Collider Spin Physics, Trieste 1995*, edited by Y. Onel, N. Paver, and A. Penzo (World Scientific Publishers, to be published), and references therein.
- [6] T.A. DeGrand and H.I. Miettinen, Phys. Rev. D **23**, 1227 (1981); **24**, 2419 (1981); **31**, 661(E) (1985); T.A. DeGrand, J. Markkanen and H.I. Miettinen, *ibid.* **32**, 2445 (1985).
- [7] B. Andersson, G. Gustafson, and G. Ingelman, Phys. Lett. **85B**, 417 (1979).
- [8] W.G.D. Dharmaratna and G.R. Goldstein, Phys. Rev. D **41**, 1731 (1990); J. Szwed, Phys. Lett. **105B**, 403 (1981); T. Fujita and T. Matsuyama, Nihon University Report No. NUP-A-879, 1987.
- [9] M. Anselmino *et al.*, Rev. Mod. Phys. **65**, 1199 (1993), and references therein.
- [10] D.P. Grosnick *et al.*, Nucl. Inst. Meth. Phys. Res. A **290**, 269 (1990).
- [11] A. Lesnik *et al.*, Phys. Rev. Lett. **35**, 770 (1975).
- [12] B.E. Bonner *et al.*, Phys. Rev. D **38**, 729 (1988).
- [13] H.T. Diehl *et al.*, Phys. Rev. Lett. **67**, 804 (1991); N.B. Wallace *et al.*, Phys. Rev. Lett. **74**, 3732 (1995).

- [14] J. Soffer and N.A. Törnqvist, Phys. Rev. Lett. **68**, 907 (1992).
- [15] R. Barni, G. Preparata, and P.G. Ratcliffe, Phys. Lett. **296B**, 251 (1992).
- [16] C. Boros and Z. Liang, Phys. Rev. D **53**, R2279 (1996); C. Boros, Z. Liang, and T. Meng, Phys. Rev. Lett. **70**, 1751 (1993); Phys. Rev. D **51**, 4867 (1995).

# TABLES

TABLE I.  $D_{NN}$  data for  $p \uparrow + p \rightarrow \Lambda^0 + X$  as a function of  $p_T$  (the errors are statistical only; systematic errors were estimated to be negligible compared to statistical ones).

$p_T$ interval (GeV/ $c$ )	$D_{NN}$	$\langle x_F \rangle$	$\langle p_T(\text{GeV}/c) \rangle$
$0.2 \leq x_F \leq 1.0$			
0.1–0.3	$-0.05 \pm 0.12$	0.42	0.23
0.3–0.5	$-0.035 \pm 0.074$	0.49	0.41
0.5–0.7	$0.147 \pm 0.071$	0.56	0.60
0.7–1.0	$0.216 \pm 0.081$	0.61	0.82
1.0–1.5	$0.26 \pm 0.17$	0.66	1.13
$0.2 \leq x_F \leq 0.5$			
0.1–0.4	$-0.03 \pm 0.10$	0.38	0.29
0.4–0.6	$0.14 \pm 0.11$	0.40	0.49
0.6–1.0	$0.20 \pm 0.13$	0.42	0.74
$0.5 \leq x_F \leq 1.0$			
0.5–0.7	$0.09 \pm 0.09$	0.65	0.60
0.7–1.0	$0.24 \pm 0.09$	0.67	0.82
1.0–1.5	$0.31 \pm 0.18$	0.69	1.14

TABLE II.  $D_{NN}$  data for  $p \uparrow + p \rightarrow \Lambda^0 + X$  as a function of  $x_F$  (the errors are statistical only).

$x_F$ interval	$D_{NN}$	$\langle x_F \rangle$	$\langle p_T (\text{GeV}/c) \rangle$
$0.1 \leq p_T \leq 1.5 \text{ GeV}/c$			
0.20–0.35	$0.03 \pm 0.13$	0.30	0.41
0.35–0.45	$0.039 \pm 0.093$	0.40	0.49
0.45–0.55	$0.079 \pm 0.082$	0.50	0.57
0.55–0.65	$0.081 \pm 0.085$	0.60	0.64
0.65–0.80	$0.148 \pm 0.086$	0.71	0.71
0.80–1.0	$0.35 \pm 0.16$	0.85	0.79
$0.1 \leq p_T \leq 0.6 \text{ GeV}/c$			
0.2–0.4	$-0.05 \pm 0.11$	0.33	0.37
0.4–0.6	$0.01 \pm 0.08$	0.49	0.42
0.6–0.8	$-0.02 \pm 0.11$	0.68	0.47
$0.6 \leq p_T \leq 1.5 \text{ GeV}/c$			
0.3–0.5	$0.17 \pm 0.12$	0.43	0.77
0.5–0.7	$0.19 \pm 0.09$	0.60	0.81
0.7–1.0	$0.37 \pm 0.11$	0.79	0.84

# FIGURES




FIG. 1.  $D_{NN}$  data as a function of  $p_T$ .




FIG. 2.  $D_{NN}$  data as a function of  $x_F$ . Also shown are  $D_{NN}$  measurements at 18.5 GeV/ $c$  from Ref. [12] ( $\langle p_T \rangle \sim 1.0$  GeV/ $c$  for the plotted points).






FIG. 3.  $D_{NN}$  data as a function of  $x_F$  divided into two  $p_T$  intervals of  $0.1 \leq p_T \leq 0.6$  GeV/ $c$  (open squares) and  $0.6 \leq p_T \leq 1.5$  GeV/ $c$  (full squares).